Proceedings of the Seventh Meeting on CPT and Lorentz Symmetry (CPT'16), Indiana University, Bloomington, June 20-24, 2016

arXiv:1607.08546v1 [hep-ph] 28 Jul 2016

General Electromagnetic Nonminimal Couplings in the Dirac Equation

J.B. Araujo, Rodolfo Casana, and M.M. Ferreira, Jr.

Departamento de Física, Universidade Federal do Maranhão
Campus Universitário do Bacanga, São Luís, Maranhão 65080-805, Brazil

We examine a new class of CPT-even and dimension-five nonminimal interactions between fermions and photons, deprived of higher-order derivatives, yielding electric dipole moment and magnetic dipole moment in the context of the Dirac equation. These couplings are Lorentz-violating nonminimal structures, composed of a rank-2 tensor, the electromagnetic tensor, and gamma matrices, being addressed in its axial and non-axial hermitian versions. We use the electron's anomalous magnetic and electric dipole moment measurements to reach upper bounds of 1 part in 10^{11} and 10^{16} GeV^{-1}.

The Standard Model (SM) structure allows for C, P, and T violations (and combinations) as long as CPT symmetry is preserved. Among the most important tests for physics beyond the SM is the search for the electric dipole moment (EDM) of particles in connection with the strong CP problem.[1,2] The EDM nonrelativistic interaction term has the form $d(\boldsymbol{\sigma}\cdot\mathbf{E})$, with $\mathbf{E}$, $\boldsymbol{\sigma}$ and d as the electric field, the spin operator, and the EDM modulus, respectively. This interaction violates both P and T symmetries, $P(\boldsymbol{\sigma}\cdot\mathbf{E})\to-(\boldsymbol{\sigma}\cdot\mathbf{E})$, $T(\boldsymbol{\sigma}\cdot\mathbf{E})\to-(\boldsymbol{\sigma}\cdot\mathbf{E})$, but preserves C and CPT symmetries. Lorentz-violating nonminimal couplings in the context of the Dirac equation that engender EDM at tree level can be stringently constrained by EDM data.[3–5] Theories endowed with Lorentz violation are a possible route for physics beyond SM that has been intensively studied in the latest years, mainly in the context of the Standard-Model Extension (SME).[6,7] Connections between LV theories and EDM physics have been addressed in Ref. 8.

In this work, a new class of dimension-five nonminimal couplings is proposed and examined with respect to the possible generation of an EDM and magnetic dipole moment (MDM),[9] having the hermiticity as the main criterion to be observed.

Nonminimal couplings in the context of the Dirac equation can be stated by modifying the usual covariant derivative.[4,5] For the cases involving a rank-2, dimensionless, LV tensor $T_{\mu\nu}$, a first proposal is given by $D_\mu = \partial_\mu + ieA_\mu + i\lambda_1 T_{\mu\nu}F^{\nu\beta}\gamma^5\gamma_\beta$. Its hermitian version is $D_\mu = \partial_\mu + ieA_\mu - \frac{i}{2}\lambda_1\left(T_{\mu\nu}F^{\nu\beta} - T_{\beta\nu}F^{\nu}{}_{\mu}\right)\gamma_\beta\gamma^5$, which leads to the modified Dirac equation

$$\left[i\gamma^\mu\partial_\mu - e\gamma^\mu A_\mu - i\lambda_1 T_{\mu\nu}F^{\nu}{}_{\beta}\sigma^{\mu\beta}\gamma^5 - m\right]\Psi = 0, \tag{1}$$

In momentum coordinates, the Dirac equation can be rewritten as $i\partial_t\Psi = \left[\boldsymbol{\alpha}\cdot\boldsymbol{\pi} + eA_0 + m\gamma^0 + H_{\rm LV}\right]\Psi$, with $H_{\rm LV} = i\lambda_1 T_{\mu\nu}F^{\nu}{}_{\beta}\gamma^0\sigma^{\mu\beta}\gamma^5$ being the LV piece of the hamiltonian. Using $F_{0j} = E^j, F_{mn} = \epsilon_{mnp}B_p$, $\sigma^{0j} = i\alpha^j$ and $\sigma^{ij} = \epsilon_{ijk}\Sigma^k$, the nonrelativistic hamiltonian for uniform fields is

$$\begin{aligned} H_{\rm NR} = H_{\rm Pauli} &+ \frac{1}{2m}\left[(\boldsymbol{\sigma}\cdot\boldsymbol{\pi})H_{12} - H_{12}(\boldsymbol{\sigma}\cdot\boldsymbol{\pi})\right] \\ &- \lambda_1\left(T_{00}\boldsymbol{\sigma}\cdot\boldsymbol{E} + T_{0i}(\boldsymbol{\sigma}\times\boldsymbol{B})^i - T_{ij}E^j\sigma^i\right), \end{aligned} \tag{2}$$

with $H_{\rm Pauli} = \frac{1}{2m}\left[\boldsymbol{\pi}^2 - e\boldsymbol{\sigma}\cdot\boldsymbol{B}\right] + eA_0$, $H_{12} = i\lambda_1(-\epsilon_{ijk}T_{i0}\sigma^k E^j - T_{ii}\left(\boldsymbol{\sigma}\cdot\boldsymbol{B}\right) + T_{ij}\sigma^j B^i)$. We notice that the terms $\lambda_1 T_{00}(\boldsymbol{\sigma}\cdot\boldsymbol{E})$, $\lambda_1 T_{ij}E^j\sigma^i$ are able to generate an EDM for the electron at tree level, so that the LV coefficients $\lambda_1 T_{00}$, $\lambda_1 T_{ij}$ can be constrained by using EDM experimental data.[10] The generation of MDM is unusual and associated with the term, $\lambda_1 T_{0i}(\sigma\times B)^i$, by an unconventional interpretation: if we consider $\lambda_1 T_{0i}(\sigma\times B)^i = \lambda_1\tilde{\boldsymbol{\sigma}}\cdot\mathbf{B} = \lambda_1\tilde{\sigma}^k B^k$, where $\tilde{\sigma}^k = \epsilon_{ijk}T_{0i}\sigma^j$ is a kind of "rotated" spin operator by the LV background, we should be able to use the MDM experimental data[11] to impose bounds on the parameter $\lambda_1 T_{0i}$ as well (if a suitable experimental set up able to measure MDM in an orthogonal direction to the applied field is found). Table I displays the obtained bounds for this coupling, its non-axial version, and other relevant cases.

Table 1. List of the bounds (in GeV^{-1} units) for the relevant couplings.

Coupling	EDM	MDM
$i\lambda_1\bar{\Psi}T_{\mu\nu}F^{\nu}{}_{\beta}\sigma^{\mu\beta}\gamma^5\Psi$	$\lvert\lambda_1 T_{00}\rvert \leq 3.8\times10^{-16}$ $\lvert\lambda_1 T\rvert \leq 1.1\times10^{-15}$	$\lvert\lambda_1 T_{0i}\rvert \leq 5.5\times10^{-11}$
$\lambda_1'\bar{\Psi}T_{\mu\nu}F^{\nu}{}_{\beta}\sigma^{\mu\beta}\Psi$	$\lvert\lambda_1' T_{i0}\rvert \leq 3.8\times10^{-16}$	$\lvert\lambda_1' T\rvert \leq 3.5\times10^{-11}$
$\lambda_4\bar{\Psi}(T_{\alpha\nu}F_{\mu\beta} - T_{\mu\beta}F_{\alpha\nu})\sigma^{\mu\beta}\sigma^{\alpha\nu}\Psi$	$\lvert\lambda_4 T_{0i}\rvert \leq 3.8\times10^{-16}$	$\lvert\lambda_4 T_{ij}\rvert \leq 5.5\times10^{-11}$

The non-axial version of this coupling is given by the following extension to the covariant derivative: $D_\mu = \partial_\mu + ieA_\mu + \frac{\lambda_1'}{2}\left(T_{\mu\nu}F^{\nu}{}_{\beta} - T_{\beta\nu}F^{\nu}{}_{\mu}\right)\gamma^\beta$, which yields a hermitian contribution to the Dirac equation,

$$\left[i\gamma^\mu\partial_\mu - e\gamma^\mu A_\mu + \lambda_1 T_{\mu\nu}F^{\nu}{}_{\beta}\sigma^{\mu\beta} - m\right]\Psi = 0, \tag{3}$$

implying the hamiltonian LV piece, $H'_{\rm LV} = -\lambda'_1 T_{\mu\nu} F^{\nu}{}_{\beta} \gamma^0 \sigma^{\mu\beta}$. The full nonrelativistic hamiltonian is

$$H_{\rm NR} = H_{\rm Pauli} + \frac{1}{2m}[(\boldsymbol{\sigma}\cdot\boldsymbol{\pi})\, H'_{12} - H'_{12}\,(\boldsymbol{\sigma}\cdot\boldsymbol{\pi})] + \lambda'_1 T_{i0}\,(\boldsymbol{\sigma}\times\boldsymbol{E})^i - \lambda'_1 T\,(\boldsymbol{\sigma}\cdot\boldsymbol{B}) + \lambda'_1 T_{ij}\sigma^j B^i, \tag{4}$$

where $T = {\rm Tr}(T_{ij})$ and $H'_{12} = i\lambda'_1(T_{00}E^i\sigma^i - T_{ij}E^j\sigma^i + T_{0i}\epsilon_{ijk}\sigma^j B^k)$. Clearly, the terms $T\,(\boldsymbol{\sigma}\cdot\boldsymbol{B})$ and $T_{ij}\sigma^j B^i$ are MDM contributions, involving only the symmetric part of the tensor $T_{\mu\nu}$, whose magnitude can be limited by the known experimental error in the MDM.[11] There is also an unconventional EDM contribution. Analogously to the magnetic case, the term $\lambda'_1 T_{i0}\,(\boldsymbol{\sigma}\times\boldsymbol{E})^i = \lambda'_1\tilde{\sigma}^k E^k$ could be considered as EDM if we take $\tilde{\sigma}^k = \epsilon_{ijk}T_{0i}\sigma^j$ as a kind of "rotated" EDM. Therefore, EDM and MDM experimental data can be used to constrain these terms, as it is shown in Table I. An interesting discussion concerning the overlapping of some bounds with others in the literature is found in Ref. 9. For a discussion on the sidereal variations, to which the LV parameters are subject, see Refs. 5, 7, 9.

The nonminimal couplings of Eqs. (1) and (3) enclose two gamma matrices contracted with the background and the $F^{\alpha\beta}$ tensor. New possibilities of couplings arise when the tensors $T_{\mu\nu}$ and $F^{\alpha\beta}$ have no mutually contracted indices, leaving three indices to be contracted with gamma matrices. These derivatives are comprised in the general expression $D_\mu = \partial_\mu + ieA_\mu + i\lambda_3 T_{\{\alpha\nu}F_{\mu\beta\}}\gamma^{\{\beta}\gamma^{\alpha}\gamma^{\nu\}}$, with the symbol $\{\}$ denoting possible permutation of the indexes μ,ν,α,β. Among them there is

$$D_\mu = \partial_\mu + ieA_\mu + i\frac{\lambda_4}{8}\,(T_{\alpha\nu}F_{\mu\beta} - T_{\mu\beta}F_{\alpha\nu})\,\gamma^\beta\sigma^{\alpha\nu}. \tag{5}$$

This coupling is distinct from the previous ones, leading to the Dirac equation contribution: $[\lambda_4 T^{\nu}{}_{\nu}F_{\mu\beta}\sigma^{\mu\beta} + i\lambda_4\,(T_{\alpha\nu}F_{\mu\beta} - T_{\mu\beta}F_{\alpha\nu})\,\sigma^{\mu\beta}\sigma^{\alpha\nu}]/8$. In this case, the symmetric part of $T_{\mu\nu}$ yields the usual MDM interaction, $\lambda_4 T^{\nu}{}_{\nu}F_{\mu\beta}\bar{\Psi}\sigma^{\mu\beta}\Psi$. Taking an antisymmetric $T_{\mu\nu}$, the modified Dirac equation assumes the form $i\partial_t\Psi = \left[\boldsymbol{\alpha}\cdot\boldsymbol{\pi} + eA_0 + m\gamma^0 + H_{\rm LV4}\right]\Psi$, where the LV piece is $H_{\rm LV4} = i\frac{\lambda_4}{8}\,[T_{\alpha\nu}F_{\mu\beta} - T_{\mu\beta}F_{\alpha\nu}]\,\gamma^0\sigma^{\mu\beta}\sigma^{\alpha\nu}$. The EDM and MDM nonrelativistic contributions are

$$H_{\rm NR(4)} = \lambda_4 T_{0i}E^j\epsilon_{ijk}\sigma^k + \lambda_4 T_{ad}B^a\sigma^d. \tag{6}$$

In this case, there are unconventional MDM and EDM contributions. The bounds achieved for this model are also included in Table I. It is possible to

show that many other couplings involving four gamma matrices are either physically redundant or do not generate spin interactions.[9]

In this work, we analyzed a new class of dimension-five, CPT-even LV nonminimal couplings between fermions and photons, composed of a general tensor, $T_{\mu\nu}$, in the context of the Dirac equation, addressing its axial and non-axial versions. EDM and MDM experimental data were used to impose upper bounds on the LV parameters of 1 part in 10^{11} and 10^{16} GeV^{-1}. The "rotated" MDM and EDM unconventional interpretation could allow constraining off-diagonal components of the LV tensor if experiments able to measure EDM and MDM components orthogonal to the applied fields are conceived.

Acknowledgments

The authors are grateful to the Brazilian research agencies CNPq, CAPES, and FAPEMA for invaluable financial support.

References

1. Y.V. Stadnik and V.V. Flambaum, Phys. Rev. D **89**, 043522 (2014); M. Pospelov and A. Ritz, Ann. Phys. **318**, 119 (2005).
2. B.L. Roberts and W.J. Marciano, *Lepton Dipole Moments*, World Scientific, Singapore, 2010.
3. P.A. Bolokhov, M. Pospelov, and M. Romalis, Phys. Rev. D **78**, 057702 (2008).
4. R. Casana, M.M. Ferreira, Jr., E. Passos, F.E.P. dos Santos, and E.O. Silva, Phys. Rev. D **87**, 047701 (2013).
5. J.B. Araujo, R. Casana, and M.M. Ferreira, Jr., Phys. Rev. D **92**, 025049 (2015).
6. D. Colladay and V.A. Kostelecký, Phys. Rev. D **55**, 6760 (1997); Phys. Rev. D **58**, 116002 (1998).
7. R. Bluhm, V.A. Kostelecký, C.D. Lane, and N. Russell, Phys. Rev. Lett. **88**, 090801 (2002); Phys. Rev. D **68**, 125008 (2003); V.A. Kostelecký and M. Mewes, Phys. Rev. D **66**, 056005 (2002).
8. I. Altarev *et al.*, Europhys. Lett. **92**, 51001 (2010); M. Haghighat, I. Motie, and Z. Rezaei, Int. J. Mod. Phys. A **28**, 1350115 (2013); A.H. Gomes, V.A. Kostelecky, and A.J. Vargas, Phys. Rev. D **90**, 076009 (2014); Y.M.P. Gomes and P.C. Malta, Phys. Rev. D **94**, 025031 (2016).
9. J.B. Araujo, R. Casana, and M.M. Ferreira, Jr., Phys. Lett. B **760**, 302 (2016).
10. J. Baron *et al.*, Science **343**, 269 (2014).
11. G. Gabrielse *et al.*, Phys. Rev. Lett. **97**, 030802 (2006); D. Hanneke, S. Fogwell, and G. Gabrielse, Phys. Rev. Lett. **100**, 120801 (2008).